

## Supplement to the paper “Asymptotic properties of the Bayes modal estimators of item parameters in item response theory”

Haruhiko Ogasawara

This note is to supplement Ogasawara (2013), and gives asymptotic expansions for the reciprocals of the estimated asymptotic standard errors used for studentized estimators and a simplification of Theorem 6.

(a)  $t$

$$\begin{aligned}
 & \text{Define } \hat{\mathbf{i}}_W^{\alpha\alpha} = (\hat{\mathbf{I}}_W^{-1})_{(\alpha\alpha)}. \text{ Then, for } t = N^{1/2}(\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0) / (\hat{\mathbf{i}}_W^{\alpha\alpha})^{1/2}, \\
 & (\hat{\mathbf{i}}_W^{\alpha\alpha})^{-1/2} = (\mathbf{i}_0^{\alpha\alpha})^{-1/2} + \frac{1}{2}(\mathbf{i}_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} \frac{2 - \delta_{ab}}{2} \{\mathbf{I}_0^{-1}(\mathbf{E}_{ab} + \mathbf{E}_{ba})\mathbf{I}_0^{-1}\}_{(\alpha\alpha)} \frac{\partial \mathbf{i}_{0ab}}{\partial \boldsymbol{\alpha}_0} (\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0) \\
 & \quad + \frac{3}{8}(\mathbf{i}_0^{\alpha\alpha})^{-5/2} \left[ \sum_{a \geq b} \frac{2 - \delta_{ab}}{2} \{\mathbf{I}_0^{-1}(\mathbf{E}_{ab} + \mathbf{E}_{ba})\mathbf{I}_0^{-1}\}_{(\alpha\alpha)} \frac{\partial \mathbf{i}_{0ab}}{\partial \boldsymbol{\alpha}_0} (\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0) \right]^2 \\
 & \quad - \frac{1}{4}(\mathbf{i}_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} \sum_{c \geq d} \frac{1}{4} (2 - \delta_{ab})(2 - \delta_{cd}) \\
 & \quad \quad \times 2 \{\mathbf{I}_0^{-1}(\mathbf{E}_{ab} + \mathbf{E}_{ba})\mathbf{I}_0^{-1}(\mathbf{E}_{cd} + \mathbf{E}_{dc})\mathbf{I}_0^{-1}\}_{(\alpha\alpha)} \frac{\partial \mathbf{i}_{0ab}}{\partial \boldsymbol{\alpha}_0} (\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0) \frac{\partial \mathbf{i}_{0cd}}{\partial \boldsymbol{\alpha}_0} (\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0) \\
 & \quad + \frac{1}{4}(\mathbf{i}_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} \frac{2 - \delta_{ab}}{2} \{\mathbf{I}_0^{-1}(\mathbf{E}_{ab} + \mathbf{E}_{ba})\mathbf{I}_0^{-1}\}_{(\alpha\alpha)} \frac{\partial^2 \mathbf{i}_{0ab}}{(\partial \boldsymbol{\alpha}_0')^{<2>}} (\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0)^{<2>} \\
 & \quad + O_p(N^{-3/2}) \\
 & \equiv (\mathbf{i}_0^{\alpha\alpha})^{-1/2} + \mathbf{i}_0^{(1)} (\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0) + \mathbf{i}_0^{(2)} (\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0)^{<2>} + O_p(N^{-3/2}) \\
 & = (\mathbf{i}_0^{\alpha\alpha})^{-1/2} + \mathbf{i}_0^{(1)} (N^{-1} \boldsymbol{\eta}_0 + N^{-1/2} \boldsymbol{\Lambda}^{(1)} \mathbf{m}^{(1)} + N^{-1} \boldsymbol{\Lambda}^{(2)} \mathbf{m}^{(2)}) \\
 & \quad + \mathbf{i}_0^{(2)} (N^{-1/2} \boldsymbol{\Lambda}^{(1)} \mathbf{m}^{(1)})^{<2>} + O_p(N^{-3/2})
 \end{aligned} \tag{S.1}$$

$$\begin{aligned}
&= (i_0^{\alpha\alpha})^{-1/2} + N^{-1/2} \mathbf{i}_0^{(1)'} \boldsymbol{\Lambda}^{(1)} \mathbf{m}^{(1)} + N^{-1} \mathbf{i}_0^{(1)'} \boldsymbol{\eta}_0 \\
&\quad + N^{-1} \{ \mathbf{i}_0^{(1)'} \boldsymbol{\Lambda}^{(2)} \mathbf{m}^{(2)} + \mathbf{i}_0^{(2)'} (\boldsymbol{\Lambda}^{(1)} \mathbf{m}^{(1)})^{<2>} \} + O_p(N^{-3/2}) \\
&\equiv (i_0^{\alpha\alpha})^{-1/2} + N^{-1} \boldsymbol{\eta}_{t0} + \sum_{j=1}^2 N^{-j/2} \boldsymbol{\lambda}_t^{(j)'} \mathbf{m}^{(j)} + O_p(N^{-3/2}),
\end{aligned}$$

where  $\sum_{a \geq b} (\cdot) = \sum_{a=1}^q \sum_{b=1}^a (\cdot)$ ,  $\delta_{ab}$  is the Kronecker delta,  $\mathbf{E}_{ab}$  is a square matrix of an appropriate size whose  $(a, b)$ th element is 1 with other ones being zero, and  $\boldsymbol{\Lambda}^{(j)}$  ( $j=1, 2$ ) are multivariate versions of  $\boldsymbol{\lambda}^{(j)}$  ( $j=1, 2$ ), respectively (see (8.2)).

In (S.1),

$$\begin{aligned}
\mathbf{i}_0^{(1)} &= \frac{1}{2} (i_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} (2 - \delta_{ab}) i_0^{\alpha a} i_0^{\alpha b} \frac{\partial i_{0ab}}{\partial \boldsymbol{\alpha}_0}, \\
\mathbf{i}_0^{(2)} &= \frac{3}{8} (i_0^{\alpha\alpha})^{-5/2} \left[ \sum_{a \geq b} (2 - \delta_{ab}) i_0^{\alpha a} i_0^{\alpha b} \frac{\partial i_{0ab}}{\partial \boldsymbol{\alpha}_0} \right]^{<2>} \\
&\quad - \frac{1}{8} (i_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} \sum_{c \geq d} (2 - \delta_{ab})(2 - \delta_{cd}) \sum_{(abcd)}^4 i_0^{\alpha a} i_0^{\alpha b} i_0^{\alpha c} i_0^{\alpha d} \frac{\partial i_{0ab}}{\partial \boldsymbol{\alpha}_0} \otimes \frac{\partial i_{0cd}}{\partial \boldsymbol{\alpha}_0} \quad (\text{S.2}) \\
&\quad + \frac{1}{4} (i_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} (2 - \delta_{ab}) i_0^{\alpha a} i_0^{\alpha b} \frac{\partial^2 i_{0ab}}{(\partial \boldsymbol{\alpha}_0)^{<2>}},
\end{aligned}$$

$\boldsymbol{\eta}_{t0} = \mathbf{i}_0^{(1)'} \boldsymbol{\eta}_0$ ,  $\boldsymbol{\lambda}_t^{(1)'} = \mathbf{i}_0^{(1)'} \boldsymbol{\Lambda}^{(1)}$ ,  $\boldsymbol{\lambda}_t^{(2)'} = \mathbf{i}_0^{(1)'} \boldsymbol{\Lambda}^{(2)} + \mathbf{i}_0^{(2)'} \boldsymbol{\Lambda}^{(1)<2>}$ , where

$\sum_{(abcd)}^4$  is the sum of four terms considering the symmetric cases for  $a, b, c$

and  $d$ , and noting  $\mathbf{I}_0 \equiv \sum_{k=1}^K \frac{1}{\pi_{0k}} \frac{\partial \pi_{0k}}{\partial \boldsymbol{\alpha}_0} \frac{\partial \pi_{0k}}{\partial \boldsymbol{\alpha}_0}$  with  $\pi_{0k} \equiv (\boldsymbol{\pi}_0)_k \equiv \{\boldsymbol{\pi}(\boldsymbol{\alpha}_0)\}_k$ , we

have

$$\begin{aligned}
\frac{\partial i_{0ab}}{\partial \alpha_{0c}} &= \sum_{k=1}^K \left( -\frac{1}{\pi_{0k}^2} \frac{\partial \pi_{0k}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0c}} + \sum_{(ab)}^2 \frac{1}{\pi_{0k}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \right), \\
\frac{\partial^2 i_{0ab}}{\partial \alpha_{0c} \partial \alpha_{0d}} &= \sum_{k=1}^K \left\{ \frac{2}{\pi_{0k}^3} \frac{\partial \pi_{0k}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0d}} \right. \\
&\quad \left. - \frac{1}{\pi_{0k}^2} \left( \sum_{(ab)}^2 \sum_{(cd)}^2 \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0d}} + \frac{\partial \pi_{0k}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0c} \partial \alpha_{0d}} \right) \right. \\
&\quad \left. + \frac{1}{\pi_{0k}} \left( \sum_{(ab)}^2 \frac{\partial^3 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c} \partial \alpha_{0d}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} + \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0b} \partial \alpha_{0d}} \right. \right. \\
&\quad \left. \left. + \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0d}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0b} \partial \alpha_{0c}} \right) \right\} \\
&(a, b, c, d = 1, \dots, q).
\end{aligned} \tag{S.3}$$

(b)  $t_H$

Using  $\hat{\mathbf{H}}_W$  (see (4.2b)), define  $\hat{h}_W^{\alpha\alpha}$ ,  $\mathbf{H}_0$ ,  $\hat{\mathbf{D}}_W$ ,  $\hat{d}_W^{\alpha\alpha}$  and  $d_0^{\alpha\alpha}$  in the following equations:

$$\begin{aligned}
t_H &\equiv \frac{N^{1/2} (\hat{\alpha}_W - \alpha_0)}{\{(-\hat{\mathbf{H}}_W^{-1})_{(\alpha\alpha)}\}^{1/2}} \equiv \frac{N^{1/2} (\hat{\alpha}_W - \alpha_0)}{(-\hat{h}_W^{\alpha\alpha})^{1/2}}, \\
-\hat{h}_W^{\alpha\alpha} &= \left[ \left\{ \sum_{k=1}^K p_k \left( -\frac{1}{\hat{\pi}_k} \frac{\partial^2 \hat{\pi}_k}{\partial \hat{\alpha}_W \partial \hat{\alpha}_W'} + \frac{1}{\hat{\pi}_k^2} \frac{\partial \hat{\pi}_k}{\partial \hat{\alpha}_W} \frac{\partial \hat{\pi}_k}{\partial \hat{\alpha}_W'} \right) \right\}^{-1} \right]_{(\alpha\alpha)}, \tag{S.4}
\end{aligned}$$

$$\hat{\mathbf{D}}_W = -\hat{\mathbf{H}}_W, \quad \mathbf{D}_0 = -\mathbf{H}_0 = \sum_{k=1}^K \pi_{Tk} \left( -\frac{1}{\pi_{0k}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_0 \partial \alpha_0'} + \frac{1}{\pi_{0k}^2} \frac{\partial \pi_{0k}}{\partial \alpha_0} \frac{\partial \pi_{0k}}{\partial \alpha_0'} \right).$$

Then,

$$\begin{aligned}
(-\hat{h}_W^{\alpha\alpha})^{-1/2} &= (\hat{d}_W^{\alpha\alpha})^{-1/2} = (d_0^{\alpha\alpha})^{-1/2} \\
&+ \frac{1}{2} (d_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} (2 - \delta_{ab}) d_0^{\alpha a} d_0^{\alpha b} \frac{\partial d_{0ab}}{\partial (\boldsymbol{\pi}_T', \boldsymbol{\alpha}_0')} \{(\mathbf{p} - \boldsymbol{\pi}_T)', (\hat{\alpha}_W - \boldsymbol{\alpha}_0)'\} \tag{S.5}
\end{aligned}$$

$$\begin{aligned}
& + \left[ \frac{3}{8} (d_0^{\alpha\alpha})^{-5/2} \left\{ \sum_{a \geq b} (2 - \delta_{ab}) d_0^{\alpha a} d_0^{\alpha b} \frac{\partial d_{0ab}}{\partial (\boldsymbol{\pi}_T', \boldsymbol{\alpha}_0')} \right\} \right]^{<2>} \\
& - \frac{1}{8} (d_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} \sum_{c \geq d^*} (2 - \delta_{ab}) (2 - \delta_{cd^*}) \\
& \quad \times \sum_{(abcd^*)}^4 d_0^{\alpha a} d_0^{bc} d_0^{d^* \alpha} \frac{\partial d_{0ab}}{\partial (\boldsymbol{\pi}_T', \boldsymbol{\alpha}_0')} \otimes \frac{\partial d_{0cd^*}}{\partial (\boldsymbol{\pi}_T', \boldsymbol{\alpha}_0')} \\
& + \frac{1}{4} (d_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} (2 - \delta_{ab}) d_0^{\alpha a} d_0^{\alpha b} \frac{\partial^2 d_{0ab}}{\{\partial (\boldsymbol{\pi}_T', \boldsymbol{\alpha}_0')\}^{<2>}} \left] \{(\mathbf{p} - \boldsymbol{\pi}_T)', (\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0)'\}^{<2>} \\
& \equiv (d_0^{\alpha\alpha})^{-1/2} + \mathbf{d}_0^{(1)} \{(\mathbf{p} - \boldsymbol{\pi}_T)', (\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0)'\} \\
& \quad + \mathbf{d}_0^{(2)} \{(\mathbf{p} - \boldsymbol{\pi}_T)', (\hat{\boldsymbol{\alpha}}_W - \boldsymbol{\alpha}_0)'\}^{<2>} + O_p(N^{-3/2}) \\
& = (d_0^{\alpha\alpha})^{-1/2} + \mathbf{d}_0^{(1)} \{N^{-1/2} \mathbf{m}^{(1)}', (N^{-1} \boldsymbol{\eta}_0 + N^{-1/2} \boldsymbol{\Lambda}^{(1)} \mathbf{m}^{(1)} + N^{-1} \boldsymbol{\Lambda}^{(2)} \mathbf{m}^{(2)})'\} \\
& \quad + \mathbf{d}_0^{(2)} \{N^{-1/2} \mathbf{m}^{(1)}', N^{-1/2} (\boldsymbol{\Lambda}^{(1)} \mathbf{m}^{(1)})'\}^{<2>} + O_p(N^{-3/2}).
\end{aligned}$$

Decomposing  $\mathbf{d}_0^{(j)}$  ( $j = 1, 2$ ) using subvectors as

$$\mathbf{d}_0^{(1)} = (\mathbf{d}_0^{(1A)}', \mathbf{d}_0^{(1B)}')' \quad \text{and} \quad \mathbf{d}_0^{(2)} = (\mathbf{d}_0^{(2AA)}', \mathbf{d}_0^{(2AB)}', \mathbf{d}_0^{(2BA)}', \mathbf{d}_0^{(2BB)}')', \quad (\text{S.6})$$

(S.5) becomes

$$\begin{aligned}
& = (d_0^{\alpha\alpha})^{-1/2} + N^{-1/2} (\mathbf{d}_0^{(1A)}' + \mathbf{d}_0^{(1B)}' \boldsymbol{\Lambda}^{(1)}) \mathbf{m}^{(1)} + N^{-1} \mathbf{d}_0^{(1B)}' \boldsymbol{\eta}_0 \\
& + N^{-1} \{ \mathbf{d}_0^{(1B)}' \boldsymbol{\Lambda}^{(2)} + \mathbf{d}_0^{(2AA)}' + 2\mathbf{d}_0^{(2AB)}' (\mathbf{I}_{(K)} \otimes \boldsymbol{\Lambda}^{(1)}) + \mathbf{d}_0^{(2BB)}' \boldsymbol{\Lambda}^{(1)<2>} \} \mathbf{m}^{(1)<2>} \\
& + O_p(N^{-3/2}) \\
& \equiv (d_0^{\alpha\alpha})^{-1/2} + N^{-1} \boldsymbol{\eta}_{H0} + \sum_{j=1}^2 N^{-j/2} \boldsymbol{\lambda}_H^{(j)}' \mathbf{m}^{(j)} + O_p(N^{-3/2}),
\end{aligned} \quad (\text{S.7})$$

where

$$\begin{aligned}
\boldsymbol{\eta}_{H0} & = \mathbf{d}_0^{(1B)}' \boldsymbol{\eta}_0, \quad \boldsymbol{\lambda}_H^{(1)}' = \mathbf{d}_0^{(1A)}' + \mathbf{d}_0^{(1B)}' \boldsymbol{\Lambda}^{(1)}, \\
\boldsymbol{\lambda}_H^{(2)}' & = \mathbf{d}_0^{(1B)}' \boldsymbol{\Lambda}^{(2)} + \mathbf{d}_0^{(2AA)}' + 2\mathbf{d}_0^{(2AB)}' (\mathbf{I}_{(K)} \otimes \boldsymbol{\Lambda}^{(1)}) + \mathbf{d}_0^{(2BB)}' \boldsymbol{\Lambda}^{(1)<2>}
\end{aligned} \quad (\text{S.8})$$

and  $\mathbf{I}_{(K)}$  is the  $K \times K$  identity matrix.

In (S.5),

$$\begin{aligned}
\frac{\partial d_{0ab}}{\partial \pi_{\text{Tk}^*}} &= -\frac{1}{\pi_{0k^*}} \frac{\partial^2 \pi_{0k^*}}{\partial \alpha_{0a} \partial \alpha_{0b}} + \frac{1}{\pi_{0k^*}^2} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0b}}, \\
\frac{\partial d_{0ab}}{\partial \alpha_{0c}} &= \sum_{k=1}^K \pi_{\text{Tk}} \left( -\frac{2}{\pi_{0k}^3} \frac{\partial \pi_{0k}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0c}} + \frac{1}{\pi_{0k}^2} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0c}} \right. \\
&\quad \left. + \sum_{(ab)}^2 \frac{1}{\pi_{0k}^2} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} - \frac{1}{\pi_{0k}} \frac{\partial^3 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0b} \partial \alpha_{0c}} \right), \\
\frac{\partial^2 d_{0ab}}{\partial \pi_{\text{Tk}^*} \partial \pi_{\text{Tl}}} &= 0, \\
\frac{\partial^2 d_{0ab}}{\partial \pi_{\text{Tk}^*} \partial \alpha_{0c}} &= -\frac{2}{\pi_{0k^*}^3} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0c}} + \frac{1}{\pi_{0k^*}^2} \frac{\partial^2 \pi_{0k^*}}{\partial \alpha_{0a} \partial \alpha_{0b}} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0c}} \\
&\quad + \frac{1}{\pi_{0k^*}^2} \sum_{(ab)}^2 \frac{\partial^2 \pi_{0k^*}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0b}} - \frac{1}{\pi_{0k^*}} \frac{\partial^3 \pi_{0k^*}}{\partial \alpha_{0a} \partial \alpha_{0b} \partial \alpha_{0c}}, \\
\frac{\partial^2 d_{0ab}}{\partial \alpha_{0c} \partial \alpha_{0d^*}} &= \sum_{k=1}^K \pi_{\text{Tk}} \left\{ \frac{6}{\pi_{0k}^4} \frac{\partial \pi_{0k}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0d^*}} \right. \\
&\quad - \frac{2}{\pi_{0k}^3} \left( \sum_{(abcd^*)}^4 \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0d^*}} + \frac{\partial \pi_{0k}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0c} \partial \alpha_{0d^*}} \right. \\
&\quad \left. \left. + \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0d^*}} \right) \right. \\
&\quad \left. + \frac{1}{\pi_{0k}^2} \left( \sum_{(abcd^*)}^4 \frac{\partial^3 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0b} \partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0d^*}} + \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0b}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0c} \partial \alpha_{0d^*}} \right. \right. \\
&\quad \left. \left. + \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0b} \partial \alpha_{0d^*}} + \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0d^*}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0b} \partial \alpha_{0c}} \right) \right. \\
&\quad \left. - \frac{1}{\pi_{0k}} \frac{\partial^4 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0b} \partial \alpha_{0c} \partial \alpha_{0d^*}} \right\} \\
&\quad (a, b, c, d^* = 1, \dots, q; k^*, l = 1, \dots, K).
\end{aligned} \tag{S.9}$$

(c)  $t_G$

Using  $\hat{\mathbf{G}}_W$  (see (4.2c)), define  $\hat{g}_W^{\alpha\alpha}$ ,  $\mathbf{G}_0$  and  $g_0^{\alpha\alpha}$  as

$$t_G \equiv \frac{N^{1/2}(\hat{\alpha}_W - \alpha_0)}{\{(\hat{\mathbf{G}}_W^{-1})_{(\alpha\alpha)}\}^{1/2}} \equiv \frac{N^{1/2}(\hat{\alpha}_W - \alpha_0)}{(\hat{g}_W^{\alpha\alpha})^{1/2}},$$

$$\mathbf{G}_0 = \sum_{k=1}^K \frac{\pi_{Tk}}{\pi_{0k}^2} \frac{\partial \pi_{0k}}{\partial \mathbf{a}_0} \frac{\partial \pi_{0k}}{\partial \mathbf{a}_0}, \quad (\text{S.10})$$

and  $g_0^{\alpha\alpha} = (\mathbf{G}_0^{-1})_{(\alpha\alpha)}$ .

Then,

$$\begin{aligned} & (\hat{g}_W^{\alpha\alpha})^{-1/2} = (g_0^{\alpha\alpha})^{-1/2} \\ & + \frac{1}{2} (g_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} (2 - \delta_{ab}) g_0^{\alpha a} g_0^{\alpha b} \frac{\partial g_{0ab}}{\partial (\boldsymbol{\pi}_T', \mathbf{a}_0')} \{(\mathbf{p} - \boldsymbol{\pi}_T)', (\hat{\mathbf{a}}_W - \mathbf{a}_0)'\}' \quad (\text{S.11}) \\ & + \left[ \frac{3}{8} (g_0^{\alpha\alpha})^{-5/2} \left\{ \sum_{a \geq b} (2 - \delta_{ab}) g_0^{\alpha a} g_0^{\alpha b} \frac{\partial g_{0ab}}{\partial (\boldsymbol{\pi}_T', \mathbf{a}_0')} \right\}^{<2>} \right. \\ & - \frac{1}{8} (g_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} \sum_{c \geq d^*} (2 - \delta_{ab})(2 - \delta_{cd^*}) \\ & \quad \times \sum_{(abcd^*)}^4 g_0^{\alpha a} g_0^{bc} g_0^{d^* \alpha} \frac{\partial g_{0ab}}{\partial (\boldsymbol{\pi}_T', \mathbf{a}_0')} \otimes \frac{\partial g_{0cd^*}}{\partial (\boldsymbol{\pi}_T', \mathbf{a}_0')} \\ & \left. + \frac{1}{4} (g_0^{\alpha\alpha})^{-3/2} \sum_{a \geq b} (2 - \delta_{ab}) g_0^{\alpha a} g_0^{\alpha b} \frac{\partial^2 g_{0ab}}{\{\partial (\boldsymbol{\pi}_T', \mathbf{a}_0')\}^{<2>}} \right] \{(\mathbf{p} - \boldsymbol{\pi}_T)', (\hat{\mathbf{a}}_W - \mathbf{a}_0)'\}'^{<2>} \\ & \equiv (g_0^{\alpha\alpha})^{-1/2} + \mathbf{g}_0^{(1)} \{(\mathbf{p} - \boldsymbol{\pi}_T)', (\hat{\mathbf{a}}_W - \mathbf{a}_0)'\}' \\ & \quad + \mathbf{g}_0^{(2)} \{(\mathbf{p} - \boldsymbol{\pi}_T)', (\hat{\mathbf{a}}_W - \mathbf{a}_0)'\}'^{<2>} + O_p(N^{-3/2}) \\ & = (g_0^{\alpha\alpha})^{-1/2} + \mathbf{g}_0^{(1)} \{N^{-1/2} \mathbf{m}^{(1)}', (N^{-1} \boldsymbol{\eta}_0 + N^{-1/2} \boldsymbol{\Lambda}^{(1)} \mathbf{m}^{(1)} + N^{-1} \boldsymbol{\Lambda}^{(2)} \mathbf{m}^{(2)})'\}' \\ & \quad + \mathbf{g}_0^{(2)} \{N^{-1/2} \mathbf{m}^{(1)}', N^{-1/2} (\boldsymbol{\Lambda}^{(1)} \mathbf{m}^{(1)})'\}'^{<2>} + O_p(N^{-3/2}). \end{aligned}$$

Decomposing  $\mathbf{g}_0^{(j)}$  ( $j = 1, 2$ ) using subvectors as

$$\mathbf{g}_0^{(1)} = (\mathbf{g}_0^{(1A)}', \mathbf{g}_0^{(1B)}')' \quad \text{and} \quad \mathbf{g}_0^{(2)} = (\mathbf{g}_0^{(2AA)}', \mathbf{g}_0^{(2AB)}', \mathbf{g}_0^{(2BA)}', \mathbf{g}_0^{(2BB)}')', \quad (\text{S.11})$$

becomes

$$\begin{aligned}
&= (\mathbf{g}_0^{\alpha\alpha})^{-1/2} + N^{-1/2} (\mathbf{g}_0^{(1A)'} + \mathbf{g}_0^{(1B)'} \Lambda^{(1)}) \mathbf{m}^{(1)} + N^{-1} \mathbf{g}_0^{(1B)'} \boldsymbol{\eta}_0 \\
&+ N^{-1} \{ \mathbf{g}_0^{(1B)'} \Lambda^{(2)} + \mathbf{g}_0^{(2AA)'} + 2\mathbf{g}_0^{(2AB)'} (\mathbf{I}_{(K)} \otimes \Lambda^{(1)}) + \mathbf{g}_0^{(2BB)'} \Lambda^{(1)\langle 2 \rangle} \} \mathbf{m}^{(1)\langle 2 \rangle} \\
&+ O_p(N^{-3/2}) \\
&\equiv (\mathbf{g}_0^{\alpha\alpha})^{-1/2} + N^{-1} \eta_{G0} + \sum_{j=1}^2 N^{-j/2} \boldsymbol{\lambda}_G^{(j)'} \mathbf{m}^{(j)} + O_p(N^{-3/2}),
\end{aligned} \tag{S.12}$$

where

$$\begin{aligned}
\eta_{G0} &= \mathbf{g}_0^{(1B)'} \boldsymbol{\eta}_0, \quad \boldsymbol{\lambda}_G^{(1)'} = \mathbf{g}_0^{(1A)'} + \mathbf{g}_0^{(1B)'} \Lambda^{(1)}, \\
\boldsymbol{\lambda}_G^{(2)'} &= \mathbf{g}_0^{(1B)'} \Lambda^{(2)} + \mathbf{g}_0^{(2AA)'} + 2\mathbf{g}_0^{(2AB)'} (\mathbf{I}_{(K)} \otimes \Lambda^{(1)}) + \mathbf{g}_0^{(2BB)'} \Lambda^{(1)\langle 2 \rangle}
\end{aligned} \tag{S.13}$$

In (S.11),

$$\begin{aligned}
\frac{\partial \mathbf{g}_{0ab}}{\partial \pi_{\text{Tk}^*}} &= \frac{1}{\pi_{0k^*}^2} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0b}}, \\
\frac{\partial \mathbf{g}_{0ab}}{\partial \alpha_{0c}} &= \sum_{k=1}^K \pi_{\text{Tk}} \left( -\frac{2}{\pi_{0k}^3} \frac{\partial \pi_{0k}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0c}} + \frac{1}{\pi_{0k}^2} \sum_{(ab)}^2 \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \right), \\
\frac{\partial^2 \mathbf{g}_{0ab}}{\partial \pi_{\text{Tk}^*} \partial \pi_{\text{Tl}}} &= 0, \\
\frac{\partial^2 \mathbf{g}_{0ab}}{\partial \pi_{\text{Tk}^*} \partial \alpha_{0c}} &= -\frac{2}{\pi_{0k^*}^3} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0c}} + \frac{1}{\pi_{0k^*}^2} \sum_{(ab)}^2 \frac{\partial^2 \pi_{0k^*}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial \pi_{0k^*}}{\partial \alpha_{0b}}, \\
\frac{\partial^2 \mathbf{g}_{0ab}}{\partial \alpha_{0c} \partial \alpha_{0d^*}} &= \sum_{k=1}^K \pi_{\text{Tk}} \left\{ \frac{6}{\pi_{0k}^4} \frac{\partial \pi_{0k}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0d^*}} \right. \\
&\quad \left. - \frac{2}{\pi_{0k}^3} \left( \sum_{(abcd^*)}^4 \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial \pi_{0k}}{\partial \alpha_{0d^*}} + \frac{\partial \pi_{0k}}{\partial \alpha_{0a}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0c} \partial \alpha_{0d^*}} \right) \right. \\
&\quad \left. + \frac{1}{\pi_{0k}^2} \left( \sum_{(ab)}^2 \frac{\partial^3 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c} \partial \alpha_{0d^*}} \frac{\partial \pi_{0k}}{\partial \alpha_{0b}} + \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0c}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0b} \partial \alpha_{0d^*}} \right. \right. \\
&\quad \left. \left. + \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0a} \partial \alpha_{0d^*}} \frac{\partial^2 \pi_{0k}}{\partial \alpha_{0b} \partial \alpha_{0c}} \right) \right\}
\end{aligned} \tag{S.14}$$

$(a, b, c, d^* = 1, \dots, q; k^*, l = 1, \dots, K)$ .

(d)  $t_R$

Define  $\hat{r}_{W\alpha\alpha}$  as

$$t_R \equiv \frac{N^{1/2}(\hat{\alpha}_W - \alpha_0)}{\left(\frac{\partial \hat{\alpha}_W}{\partial \mathbf{p}'} \hat{\mathbf{\Omega}}_T \frac{\partial \hat{\alpha}_W}{\partial \mathbf{p}}\right)^{1/2}} \equiv \frac{N^{1/2}(\hat{\alpha}_W - \alpha_0)}{\hat{r}_{W\alpha\alpha}^{1/2}}, \quad (\text{S.15})$$

where  $\frac{\partial \hat{\alpha}_W}{\partial \mathbf{p}} = \frac{\partial \alpha_W(\mathbf{p})}{\partial \mathbf{p}} \Big|_{\mathbf{p}=\mathbf{p}}$ .

$$\text{Recalling } \frac{\partial \alpha_W}{\partial \boldsymbol{\pi}_T} = \frac{\partial \alpha_W(\mathbf{p})}{\partial \mathbf{p}} \Big|_{\substack{\mathbf{a}=\mathbf{a}_W \\ \mathbf{p}=\boldsymbol{\pi}_T}} = \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T} + N^{-1} \frac{\partial \alpha_{\Delta W}}{\partial \boldsymbol{\pi}_T} + O(N^{-2}) \quad (\text{see (3.4)}),$$

and using the definitions

$$r_{W\alpha\alpha} \equiv \frac{\partial \alpha_W}{\partial \boldsymbol{\pi}_T'} \boldsymbol{\Omega}_T \frac{\partial \alpha_W}{\partial \boldsymbol{\pi}_T}, \quad r_{0\alpha\alpha} \equiv \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T'} \boldsymbol{\Omega}_T \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T} \quad (\text{S.16})$$

$$\text{and } r_{\Delta W 0} \equiv \frac{\partial \alpha_{\Delta W}}{\partial \boldsymbol{\pi}_T'} \boldsymbol{\Omega}_T \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T},$$

we have

$$\begin{aligned} \hat{r}_{W\alpha\alpha}^{-1/2} &= r_{W\alpha\alpha}^{-1/2} - \frac{1}{2} r_{W\alpha\alpha}^{-3/2} \frac{\partial r_{Wab}}{\partial \boldsymbol{\pi}_T'} (\mathbf{p} - \boldsymbol{\pi}_T) + \frac{3}{8} r_{W\alpha\alpha}^{-5/2} \left( \frac{\partial r_{Wab}}{\partial \boldsymbol{\pi}_T'} \right)^{\langle 2 \rangle} (\mathbf{p} - \boldsymbol{\pi}_T)^{\langle 2 \rangle} \\ &\quad - \frac{1}{4} r_{W\alpha\alpha}^{-3/2} \frac{\partial^2 r_{Wab}}{(\partial \boldsymbol{\pi}_T')^{\langle 2 \rangle}} (\mathbf{p} - \boldsymbol{\pi}_T)^{\langle 2 \rangle} + O_p(N^{-3/2}), \end{aligned} \quad (\text{S.17})$$

where

$$\begin{aligned} r_{W\alpha\alpha} &= r_{0\alpha\alpha} + 2N^{-1} r_{\Delta W 0} + O(N^{-2}), \\ r_{W\alpha\alpha}^{-1/2} &= r_{0\alpha\alpha}^{-1/2} - N^{-1} r_{0\alpha\alpha}^{-3/2} r_{\Delta W 0} + O(N^{-2}), \\ r_{W\alpha\alpha}^{-j/2} &= r_{0\alpha\alpha}^{-j/2} + O(N^{-1}) \quad (j=3,5), \\ \frac{\partial r_{Wab}}{\partial \boldsymbol{\pi}_T} &= \frac{\partial r_{0ab}}{\partial \boldsymbol{\pi}_T} + O(N^{-1}), \quad \frac{\partial^2 r_{Wab}}{(\partial \boldsymbol{\pi}_T)^{\langle 2 \rangle}} = \frac{\partial^2 r_{0ab}}{(\partial \boldsymbol{\pi}_T)^{\langle 2 \rangle}} + O(N^{-1}). \end{aligned} \quad (\text{S.18})$$

From the above results, (S.17) becomes



$$\begin{aligned}
&= r_{0\alpha\alpha}^{-1/2} - N^{-1} r_{0\alpha\alpha}^{-3/2} r_{\Delta W 0} - \frac{N^{-1/2}}{2} r_{0\alpha\alpha}^{-3/2} \frac{\partial r_{0ab}}{\partial \boldsymbol{\pi}_T} \mathbf{m}^{(1)} \\
&\quad + N^{-1} \left\{ \frac{3}{8} r_{0\alpha\alpha}^{-5/2} \left( \frac{\partial r_{0\alpha\alpha}}{\partial \boldsymbol{\pi}_T} \right)^{\langle 2 \rangle} - \frac{1}{4} r_{0\alpha\alpha}^{-3/2} \frac{\partial^2 r_{0\alpha\alpha}}{(\partial \boldsymbol{\pi}_T)^{\langle 2 \rangle}} \right\} \mathbf{m}^{(1)\langle 2 \rangle} + O_p(N^{-3/2}) \quad (\text{S.19})
\end{aligned}$$

$$\equiv r_{0\alpha\alpha}^{-1/2} + N^{-1} \eta_{R0} + \sum_{j=1}^2 N^{-j/2} \boldsymbol{\lambda}_R^{(j)} \mathbf{m}^{(j)} + O_p(N^{-3/2}),$$

$$\text{where } \eta_{R0} = -r_{0\alpha\alpha}^{-3/2} r_{\Delta W 0}, \quad \boldsymbol{\lambda}_R^{(1)} = -\frac{1}{2} r_{0\alpha\alpha}^{-3/2} \frac{\partial r_{0\alpha\alpha}}{\partial \boldsymbol{\pi}_T}$$

$$\text{and } \boldsymbol{\lambda}_R^{(2)} = \frac{3}{8} r_{0\alpha\alpha}^{-5/2} \left( \frac{\partial r_{0\alpha\alpha}}{\partial \boldsymbol{\pi}_T} \right)^{\langle 2 \rangle} - \frac{1}{4} r_{0\alpha\alpha}^{-3/2} \frac{\partial^2 r_{0\alpha\alpha}}{(\partial \boldsymbol{\pi}_T)^{\langle 2 \rangle}}.$$

In (S.19), using  $\frac{\partial \boldsymbol{\Omega}_T}{\partial \pi_{Tj}} = \mathbf{E}_{jj} - \boldsymbol{\pi}_T \mathbf{e}_j' - \mathbf{e}_j \boldsymbol{\pi}_T'$  where  $\mathbf{e}_j$  is the vector

whose  $j$ -th element is 1 with other ones being zero, we have

$$\begin{aligned}
\frac{\partial r_{0\alpha\alpha}}{\partial \pi_{Tj}} &= 2 \frac{\partial^2 \alpha_0}{\partial \pi_{Tj} \partial \boldsymbol{\pi}_T} \boldsymbol{\Omega}_T \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T} + \left( \frac{\partial \alpha_0}{\partial \pi_{Tj}} \right)^2 - 2 \frac{\partial \alpha_0}{\partial \pi_{Tj}} \boldsymbol{\pi}_T' \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T}, \\
\frac{\partial^2 r_{0\alpha\alpha}}{\partial \pi_{Tj} \partial \pi_{Tk}} &= 2 \frac{\partial^3 \alpha_0}{\partial \pi_{Tj} \partial \pi_{Tk} \partial \boldsymbol{\pi}_T} \boldsymbol{\Omega}_T \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T} + 2 \frac{\partial^2 \alpha_0}{\partial \pi_{Tj} \partial \boldsymbol{\pi}_T} \boldsymbol{\Omega}_T \frac{\partial^2 \alpha_0}{\partial \pi_{Tk} \partial \boldsymbol{\pi}_T} \quad (\text{S.20}) \\
&\quad + \sum_{(jk)}^2 2 \left( \frac{\partial^2 \alpha_0}{\partial \pi_{Tj} \partial \pi_{Tk}} \frac{\partial \alpha_0}{\partial \pi_{Tk}} - \frac{\partial^2 \alpha_0}{\partial \pi_{Tj} \partial \boldsymbol{\pi}_T} \boldsymbol{\pi}_T' \frac{\partial \alpha_0}{\partial \pi_{Tk}} \right) \\
&\quad - 2 \frac{\partial \alpha_0}{\partial \pi_{Tj}} \frac{\partial \alpha_0}{\partial \pi_{Tk}} - 4 \frac{\partial^2 \alpha_0}{\partial \pi_{Tj} \partial \pi_{Tk}} \boldsymbol{\pi}_T' \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T} \quad (j, k = 1, \dots, q).
\end{aligned}$$

(e)  $\eta_{B-CV0}$

The quantity  $\eta_{V0}$  is the generic expression for  $\eta_{t0}(= \mathbf{i}_0^{(1)'} \boldsymbol{\eta}_0)$ ,  $\eta_{H0}(= \mathbf{d}_0^{(1B)'} \boldsymbol{\eta}_0)$ ,  $\eta_{G0}(= \mathbf{g}_0^{(1B)'} \boldsymbol{\eta}_0)$  and  $\eta_{R0} \left( = -r_{0\alpha\alpha}^{-3/2} r_{\Delta W0} = -r_{0\alpha\alpha}^{-3/2} \frac{\partial \alpha_{\Delta W}}{\partial \boldsymbol{\pi}_T'} \boldsymbol{\Omega}_T \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T'} \right)$ . The corresponding  $\eta_{B-CV0}$  is generically defined for  $\eta_{B-Ct0}(= \mathbf{i}_0^{(1)'} \boldsymbol{\eta}_{B-C0})$ ,  $\eta_{B-CH0}(= \mathbf{d}_0^{(1B)'} \boldsymbol{\eta}_{B-C0})$ ,  $\eta_{B-CG0}(= \mathbf{g}_0^{(1B)'} \boldsymbol{\eta}_{B-C0})$  and  $\eta_{B-CR0} \left( = r_{0\alpha\alpha}^{-3/2} \frac{\partial \beta_{ML1}}{\partial \boldsymbol{\pi}_T'} \boldsymbol{\Omega}_T \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T'} \right)$ .

(f) **A simplification of Theorem 6.**

$\kappa_2(w_{B-P}^*)$  can be simplified as follows. Using (8.20), (8.19) becomes

$$\begin{aligned} \frac{\partial \boldsymbol{\alpha}_{B-P}}{\partial \boldsymbol{\pi}_T'} &= -\boldsymbol{\Lambda}^{-1} \frac{\partial^2 l_{ML}}{\partial \boldsymbol{\alpha}_0 \partial \boldsymbol{\pi}_T'} - N^{-1} \frac{\partial \boldsymbol{\beta}_{ML1}}{\partial \boldsymbol{\pi}_T'} + O(N^{-2}) \\ &\equiv \frac{\partial \boldsymbol{\alpha}_0}{\partial \boldsymbol{\pi}_T'} + N^{-1} \frac{\partial \boldsymbol{\alpha}_{\Delta B-P}}{\partial \boldsymbol{\pi}_T'} + O(N^{-2}), \end{aligned}$$

which gives

$$\begin{aligned} \kappa_2(w_{B-P}^*) &= \beta_{ML2} + N^{-1} \left( \beta_{ML\Delta 2} + 2 \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T'} \boldsymbol{\Omega}_T \frac{\partial \alpha_{\Delta B-P}}{\partial \boldsymbol{\pi}_T'} \right) + O(N^{-2}) \\ &\left( \beta_{B-P2} = \beta_{ML2}, \beta_{B-P\Delta 2} = \beta_{B-CML\Delta 2} = \beta_{ML\Delta 2} - 2 \frac{\partial \alpha_0}{\partial \boldsymbol{\pi}_T'} \boldsymbol{\Omega}_T \frac{\partial \beta_{ML1}}{\partial \boldsymbol{\pi}_T'} \right). \end{aligned}$$

That is, we find that the asymptotic cumulants of  $\hat{\alpha}_{B-P}$  up to the fourth order and the higher-order asymptotic variance are identical to those of  $\hat{\alpha}_{B-CML}$  or the bias-corrected ML estimator, respectively.

### Reference

Ogasawara, H. (2013). Asymptotic properties of the Bayes modal estimators of item parameters in item response theory. *Computational Statistics*, 28 (6), 2559-2583.